

# No short-term effect of closing a rubbish dump on reproductive parameters of an Egyptian Vulture population in Turkey

JAKOB KATZENBERGER, EVRIM TABUR, BİLGEÇAN ŞEN,  
SÜREYYA İSFENDİYAROĞLU, İTRİ LEVENT ERKOL and STEFFEN OPPEL

## Summary

Changes in food availability that lead to lower reproductive output or lower survival probability are important drivers of the widespread declines in vulture populations. Permanent feeding stations for scavengers, such as vulture restaurants or rubbish dumps, may have both positive and negative effects on reproductive parameters. Here we examine the effects of the closure of a large communal rubbish dump on breeding success and fledging rate of a dense population of the 'Endangered' Egyptian Vulture *Neophron percnopterus* in central Turkey to assess whether the closure may have affected the population. We monitored territories from 2011 to 2016, and tested whether the closure of the rubbish dump in early 2015 coincided with changes in reproductive parameters while accounting for confounding variables such as weather and the availability of other predictable foraging opportunities. We found an average productivity of 0.78 fledglings per territorial pair before the dump closed and 0.82 after the closure, an average breeding success of 0.64 before and 0.71 after the closure, and an average fledging rate of 1.17 fledglings per successful pair before and 1.26 after the closure of the rubbish dump. Once confounding variables were accounted for, the closure of the rubbish dump did not have a significant effect on reproductive parameters ( $P = 0.426$  for nest survival and  $P = 0.786$  for fledging rate). We speculate that the Egyptian Vulture population in central Turkey may have sufficient alternative food sources and high levels of intra-specific competition due to its density, so that the closure of the rubbish dump may not have resulted in detectable positive or negative effects. We recommend the maintenance of small traditional animal husbandry farms and disposal practices that mimic the spatio-temporally unpredictable supply of food sources that appears to be most beneficial for avian scavengers.

## Introduction

Among the raptor species of the world, avian scavengers like vultures have shown some of the most dramatic population declines (Thiollay 2006, Virani *et al.* 2011, Chaudhary *et al.* 2012, Ogada *et al.* 2012, 2016). One factor contributing to widespread declines of vulture populations is changes to agricultural practices and sanitary regulations that have fundamentally altered food availability and resulted in lower reproductive output or lower survival probability (Carrete *et al.* 2007, Donazar *et al.* 2010, Margalida *et al.* 2014b). Most prominently, the European Union legislation that mandated the removal of livestock carcasses from the landscape has led to food shortages and demographic changes owing to food now being provided at large, predictable feeding stations (Donazar *et al.* 2009, Margalida *et al.* 2010, Margalida and Colomer 2012). Although these feeding stations have supported many vulture populations (González *et al.* 2006, Oro *et al.* 2008,

Cortés-Avizanda *et al.* 2010), their predictable nature can attract a large number of non-breeding birds that interfere and compete with breeding birds and ultimately reduce productivity (Carrete *et al.* 2006, Cortés-Avizanda *et al.* 2010, Lieury *et al.* 2015). Understanding the influence of artificial and predictable food provisioning on the demography of populations is therefore important to inform management strategies (López-López *et al.* 2014, Margalida *et al.* 2014b).

For many vulture species, accumulating data on important demographic processes becomes logistically challenging once populations have declined and birds nest at low density over vast geographic areas. Obtaining solid baseline information while populations still nest at high density is therefore vital to inform the effects of changing food availability resulting from socio-economic or sanitary policy changes. Such changes are occurring increasingly rapidly in countries that sacrifice natural resources for the purpose of economic growth (Donald *et al.* 2006, 2015, Kamp *et al.* 2015). Turkey is a large country that connects Europe and Asia, and hosts an enormously diverse biodiversity which is under increasing threat of rapid economic expansion (Şekercioğlu *et al.* 2011). Besides hosting a vast range of resident wildlife, Turkey is also a gateway for millions of western Palaearctic migratory birds to reach wintering areas in Africa (Cameron *et al.* 1967, Porter and Willis 1968, Sutherland and Brooks 1981, Opper *et al.* 2014), and for some migratory species the populations in Turkey may support populations in Europe (Demerdzhiev *et al.* 2015). However, despite harbouring globally significant populations of many species, the demographic consequences of rapidly changing landscape structures on Turkey's biodiversity are generally poorly understood (Şekercioğlu *et al.* 2011).

One species for which Turkey holds a very important population is the Egyptian Vulture *Neophron percnopterus*, which has a European population between 3,300 and 5,050 breeding pairs mostly distributed in Spain (1,500 pairs) and Turkey (1,500–3,000 pairs) (Iñigo *et al.* 2008, Şen *et al.* 2017). The species is globally 'Endangered' due to long term declines in Europe, Africa and India (Cuthbert *et al.* 2006, Thiollay 2006, Velevski *et al.* 2015, Ogada *et al.* 2016), and these declines are mainly caused by poisoning, direct persecution, decreased food availability, electrocution and collision with power lines and wind turbines (Liberatori and Penteriani 2001, Mateo-Tomás *et al.* 2010, Angelov *et al.* 2013, Sanz-Aguilar *et al.* 2015, Velevski *et al.* 2015). The Egyptian Vulture population in Turkey is not only important due to its size, but also due to its geographic position that connects it to flyways from eastern Europe and central Asia. The declining population on the Balkan peninsula migrates mostly through Turkey (Opper *et al.* 2015), and immature Egyptian Vultures originating from the Balkans have been prospecting known breeding sites in Turkey (Bougain 2016). Safeguarding and restoring the Balkan population might therefore require a strong and productive population in Turkey, as has been shown for Egyptian Vulture populations elsewhere and for other raptors in the region (Demerdzhiev *et al.* 2015, Lieury *et al.* 2015, Tauler *et al.* 2015). Despite the importance of the Egyptian Vulture population in Turkey, population size and trends are unknown and reliable information is needed to assess the consequences of changing agricultural and land use policies on the species' status (Kirwan *et al.* 2008, Şen *et al.* 2017).

Changes in land use, away from traditional livestock farming to a more intensive agriculture, are believed to have detrimental effects on Egyptian Vultures because of reduced food availability (Mateo-Tomás and Olea 2010, Margalida *et al.* 2012, Dobrev *et al.* 2016), and this degradation of feeding habitats is speculated to pose a key threat for the species in Turkey (Iñigo *et al.* 2008). In addition, sanitary EU legislations (Regulation CE 1774/2002) that prohibit carcass disposal in the countryside, which negatively affected vultures in Spain, were adopted by the Turkish government as part of the EU accession process in March 2010 (Turkish Ministry of Environment and Urbanisation directive on sanitary landfills OG. 26.03.2010/ 27533). These new regulations may affect food availability for vultures because they limit the disposal of livestock carcasses in the landscape and mandate that landfills reduce the amount of accessible food waste. Conversely, Egyptian Vultures have very high dietary plasticity (Margalida and Colomer 2012, Margalida *et al.* 2012, Dobrev *et al.* 2016), and the removal of a predictable food source such as a rubbish dump may be easily compensated by a breeding population, or indeed have positive effects due to reduced competition and interaction with immature birds frequenting the predictable food source

(Cortés-Avizanda *et al.* 2010, Lieury *et al.* 2015). However, there is currently no quantitative information whether the removal of established food sources affects Egyptian Vultures in Turkey.

In this study we examine whether the closure of a predictable food source affected the reproductive parameters of a sizeable Egyptian Vulture breeding population in Central Anatolia, Turkey. We monitored breeding performance of a dense population from 2011 to 2016, and examined whether the closure of a central rubbish dump, which used to contain human food waste and animal carcasses and therefore functioned as a reliable food source for vultures up to 2014, resulted in a change in reproductive parameters in 2015 and 2016. Although this specific rubbish dump did not affect nest site selection of the Egyptian Vulture population (Şen *et al.* 2017), recent evidence from Spain suggests that territory occupancy may be influenced by such predictable food sources (Tauler-Ametller *et al.* 2017). This study provides the first baseline information on the breeding performance of an Egyptian Vulture population in Turkey and explores the effect of sanitary measures on reproductive parameters that have led to major changes of vulture populations in western Europe (Margalida *et al.* 2010, 2014b).

## Methods

### Study area

We studied Egyptian Vultures near the town of Beypazari in Ankara province in Central Anatolia, Turkey (40°05'N, 31°53'E; Figure 1). The study area (1,293 km<sup>2</sup>) ranges from 400 m to 1,800 m elevation, and contains high biodiversity which is captured by three Key Biodiversity Areas: Kirmir Valley, Sariyar Dam and Nallihan Hills (Eken *et al.* 2006). The southern and western parts of the study area are mainly composed of grasslands with low-intensity livestock grazing along with intensive agriculture, while the northern part is more forested and mountainous. Traditional and industrial agricultural practices provide three principal types of feeding opportunities for




Figure 1. Map of the study area Beypazari in the Ankara province in Turkey. Egyptian Vulture nest locations are displayed in relation to possible food sources.

vultures (Figure 1): sheep pens, chicken farms, and chicken farm disposal sites. While sheep pens and chicken farms occasionally discard livestock carcasses, the chicken farm disposal sites hold large quantities of chicken remains and excrement which are spread on grasslands to act as fertilizer. The single-most important congregation site for Egyptian Vultures with up to 50 birds attending simultaneously used to be the municipal rubbish dump of Beypazarı. This facility was officially closed in March 2015 because the close proximity of the rubbish dump to existing and developing settlements and recreational areas created a health hazard risk to the public. The rubbish dump closure provided the opportunity to specifically test whether the removal of such a predictable food source affected the reproductive parameters of Egyptian Vultures.

### *Field data collection and calculation of environmental variables*

Breeding attempts of Egyptian Vultures were monitored each year from 2011 to 2016, usually from pre-laying until fledging (early April until late August), but a small number of nests were only found after incubation had been initiated. We included only territories where the exact location of the nest could be confirmed. The number of monitored territories increased between 2011 and 2014 due to additive search efforts in every breeding season. Within a breeding season, each monitored territory was visited at least once per month, and we determined the number of fledglings produced by observing dependent young being fed by adults in the vicinity of the nest at the end of the breeding season (August). Observations at nest sites were made with a 20–60× magnification spotting scope at distances varying between 300 and 800 m from nesting cliffs.

We recorded all predictable food source locations (sheep pens, chicken farms and chicken farm disposal sites) using a handheld GPS device during field work and further mapped additional sites using satellite images of the study area. Because exposure to weather may affect breeding performance of raptors (Vlachos *et al.* 1998, Liberatori and Penteriani 2001, Sarà and Vittorio 2003, García-Ripollés and López-López 2006), we obtained daily weather variables (mean temperature and precipitation) in our study area for all years from the National Centres for Environmental Prediction downloaded via the R-package RNCEP (Kemp *et al.* 2012), and averaged temperature and summed precipitation for each breeding season for analysis.

Because access to food and social interactions have been found to affect Egyptian Vulture demography elsewhere (Vlachos *et al.* 1998, Liberatori and Penteriani 2001, Sarà and Vittorio 2003, García-Ripollés and López-López 2006, Carrete *et al.* 2007), we calculated several environmental variables for each monitored vulture nest that could also explain variation in reproductive parameters besides the presence of the rubbish dump. We enumerated the number of sheep pens, chicken farms, and chicken farm disposal sites within a 20-km radius around each nest, reflecting the area that is easily accessible to territorial adult vultures (López-López *et al.* 2014). To characterise intraspecific competition, we calculated the distance to the nearest active Egyptian Vulture nest, and counted the number of active nests within a 2-km radius. To test the effect of the rubbish dump closure we calculated the distance from each nest to the rubbish dump.

### *Data analysis*

Food availability, weather, and competition may affect three biological processes that comprise overall annual productivity: whether territorial pairs initiated a breeding attempt (breeding propensity), whether pairs that did initiate a breeding attempt raised any fledglings (breeding success), and whether successfully breeding pairs managed to raise two rather than just one fledgling (fledging rate). Because our monitoring was mostly initiated around nesting pairs, we here only investigate the effects of environmental variables on breeding success and fledging rate, and provide summaries of annual productivity, which we define as the number of fledglings raised by a territorial pair (including non-breeding pairs).

To test whether the rubbish dump closure in 2015 could effectively explain variation in breeding success and fledging rate, we constructed two competing models for each reproductive parameter to examine the main hypothesis of management interest, and compared these two models using a likelihood-ratio test (Lewis *et al.* 2011). Because many different factors can affect Egyptian Vulture reproductive parameters (Oppel *et al.* 2017), our basic model considered that breeding success or fledging rate could vary with the number of nearby active nests and the distance to the nearest nest (intraspecific competition); the number of sheep pens, chicken farms and chicken farm disposal sites within a 20-km radius (food availability); and total rainfall and average temperature of a breeding season (weather effects). Our competing model included all the effects of the basic model, plus the effect of the presence (2011–2014) or absence (2015–2016) of the rubbish dump. Because the effect of the rubbish dump may have depended on the distance between the nest and the rubbish dump, we specified this effect as the distance from each nest to the rubbish dump for years when the rubbish dump was available (2011–2014) and zero for years when the dump was not available (2015–2016). An alternative parameterisation using the availability of the rubbish dump as a binary variable yielded virtually identical results that are not presented here.

We used generalised linear mixed models with a binomial error distribution to evaluate these hypotheses on two different reproductive parameters (breeding success, fledging rate) while accounting for variation between territories and years by including the territory and the year as random intercepts in each model (Gillies *et al.* 2006, Bolker *et al.* 2009). We standardized all continuous variables and fitted all models using the Laplace approximation in R 3.2.5 (R Core Team 2016) with the package 'lme4' version 1.1-7 (Bates *et al.* 2015).

To account for the varying exposure time of nests in our analysis of breeding success, we used the Mayfield logistic regression (Hazler 2004), which incorporates the exposure time of each nest in an adapted link function (available at: <http://rpubs.com/bbolker/logregexp>) to address the problem that nests that fail earlier may be less likely to be accounted for during monitoring. We removed the data from 2012 for the analysis of breeding success, because in 2012 nest monitoring was carried out only in July and August.

We evaluated the explanatory power of models by predicting the response variable based on the full model and data set, and then calculated the area under the receiver-operated characteristic curve (AUC), a common performance metric for binary data that indicates whether the model has a poor (0.5) or good (1.0) discriminative ability (Fielding and Bell 1997, Jiménez-Valverde 2012).

## Results

### *Breeding population size and overall productivity*

The number of Egyptian Vulture territories monitored in the Beypazarı population increased from 37 in 2011 to 81 in 2014, but because the survey effort was not exhaustive, the actual population size may be higher. The density of Egyptian Vultures in the study area was therefore at least 6.26 pairs per 100 km<sup>2</sup>. Breeding propensity ranged between 0.83 and 0.97 with an average of 0.87 before and 0.91 after the dump closure; breeding success ranged from 0.61 to 0.76 with an average of 0.64 before and 0.71 after the closure; and annual productivity ranged from 0.71 to 0.92 fledglings/territorial pair, with an average of 0.78 before the dump closed and 0.82 after the closure (Table 1).

During the Egyptian Vulture breeding season from May to August in 2011–2016, average daily temperatures were 24°C (range: 11–34°C) and precipitation ranged from 203 to 590 mm per season. The full model explaining variation in breeding success of 244 monitored breeding attempts had a reasonable discriminative capacity (AUC = 0.707), but there was no evidence that breeding success was related to the distance from the rubbish dump in years when the dump was operational (Likelihood ratio test,  $\chi^2 = 0.634$ ,  $P = 0.426$ ). The only parameter that was estimated to be significantly different from zero indicated that nest failure rate increased with increasing distance to the nearest occupied Egyptian Vulture nest (Table 2).

Table 1. Mean ( $\pm$  95% confidence interval) reproductive parameters of the Egyptian Vulture population around Beypazarı, Turkey, during a period with an open rubbish dump (2011–2014) and after the rubbish dump had been closed (2015–2016). Note that the number of nests monitored is a consequence of survey effort and not related to changes in population size. See text for definition on how reproductive parameters were calculated.

Year	breeding propensity			breeding success			fledging rate			productivity		
	<i>n</i>	mean	95% conf. int.	<i>n</i>	mean	95% conf. int.	<i>n</i>	mean	95% conf. int.	<i>n</i>	mean	95% conf. int.
2011				37	0.70	(0.54-0.83)	26	1.19	(1.03-1.35)	40	0.84	(0.63-1.04)
2012	56	0.83*	(0.7-0.91)	40	0.90*	(0.76-0.96)	36	1.31	(1.17-1.44)	56	1.17*	(0.98-1.37)
2013	106	0.97	(0.89-0.99)	63	0.62	(0.49-0.73)	39	1.26	(1.13-1.39)	106	0.78	(0.62-0.94)
2014	168	0.85	(0.76-0.91)	69	0.61	(0.49-0.72)	42	1.17	(1.04-1.29)	168	0.71	(0.56-0.86)
2015	44	0.86	(0.73-0.94)	38	0.76	(0.6-0.87)	29	1.21	(1.05-1.36)	44	0.92	(0.72-1.12)
2016	40	0.88	(0.75-0.95)	38	0.73	(0.57-0.85)	27	1.19	(1.03-1.34)	40	0.86	(0.66-1.07)

\* nest monitoring in 2012 started in July, and may have therefore overestimated breeding success and underestimated breeding propensity because early nest failures went unrecorded.

Table 2. Parameter estimates of the fixed effects of two generalised linear mixed models evaluating the effect of the closure of the rubbish dump (in bold) while accounting for other factors affecting reproductive parameters of Egyptian Vultures around Beypazarı, Turkey, 2011–2016. See text for description of models; note that breeding success is modelled as nest failure rate. SE = standard error of the estimate.

Response variable	Parameter	Estimate	SE	<i>z</i>	<i>P</i>
nest failure rate	(Intercept)	4.03	0.27	14.64	0.00
	total rainfall over breeding season	0.08	0.11	0.68	0.50
	mean temperature over breeding season	-0.30	0.21	-1.46	0.14
	N chicken dump sites	0.14	0.21	0.65	0.52
	N sheep pens	0.05	0.23	0.22	0.82
	N chicken farms	-0.04	0.19	-0.22	0.82
	N nests	0.09	0.08	1.08	0.28
	distance to nearest occupied nest	0.16	0.07	2.40	0.02
	<b>distance to rubbish dump when available</b>	<b>-0.02</b>	<b>0.02</b>	<b>-0.80</b>	<b>0.43</b>
fledging rate	(Intercept)	-1.88	0.44	-4.32	0.00
	total rainfall over breeding season	-0.21	0.21	-1.02	0.31
	mean temperature over breeding season	0.05	0.18	0.26	0.80
	N chicken dump sites	0.72	0.45	1.61	0.11
	N sheep pens	-0.26	0.46	-0.56	0.58
	N chicken farms	0.34	0.41	0.84	0.40
	distance to nearest occupied nest	0.15	0.12	1.32	0.19
	<b>distance to rubbish dump when available</b>	<b>0.01</b>	<b>0.03</b>	<b>0.27</b>	<b>0.79</b>

The mean fledging rate of 199 successful breeding attempts was  $1.22 \pm 0.41$  (Table 1). The full model explaining variation in fledging rate had good discriminative capacity (AUC = 0.813), but there was no evidence that fledging rate was related to the distance from the rubbish dump in years when the dump was operational ( $\chi^2 = 0.074$ ,  $P = 0.786$ ). Parameter estimates of all other environmental factors had too low precision to be statistically different from zero (Table 2).

## Discussion

Our analysis of Egyptian Vulture reproductive parameters shows annual variability in breeding success but no statistical support for a distinct effect of the closure of a main food source, the communal rubbish dump. Instead, we found that Egyptian Vultures in this dense breeding population appear to have higher breeding success when nesting in close proximity to other pairs, which may be a consequence of spatially aggregated ideal nesting opportunities (Şen *et al.* 2017).

Our estimate of mean annual productivity (0.79) is slightly lower than the mean productivity of European populations (Iñigo *et al.* 2008). Across Europe, productivity ranges from 0.6 to 1.04 fledglings per territorial pair among both declining (Grubač *et al.* 2014, Oppel *et al.* 2017) and increasing populations (Mateo-Tomás *et al.* 2010, Tauler *et al.* 2015). The productivity for the Beypazarı population was very similar to the productivity of an increasing population in southern France that was aided by vulture restaurants (Lieury *et al.* 2015). However, the French population benefited from immigration and productivity suffered from compensatory density feedback possibly due to increased interference of floaters (Lieury *et al.* 2015). We found that breeding success appeared to increase when nests were closer to a neighbouring nest (Table 2), which is contrary to the expectation that interference competition with conspecifics might adversely affect productivity. However, this result might explain why the distance to the nearest nest in this population is much lower than in other European populations and that conspecific attraction appeared to peak at distances  $\sim 1,500$  m from the nearest nest (Şen *et al.* 2017). Whether this aggregated nesting is due to the spatial proximity of certain topographic or habitat attributes that Egyptian Vultures prefer for nesting, or whether nesting in close proximity actually confers a fitness advantage by some other process will require further study.



In Beypazarı, Egyptian Vultures are breeding at a density of more than six pairs per 100 km<sup>2</sup>, which is considerably higher than early records from Spain when Ceballos and Donázar (1989) found approximately 1.4 pairs/100 km<sup>2</sup>, and significantly denser than the current figures from Northern Spain where only 0.14 territories/100 km<sup>2</sup> exist (Mateo-Tomás and Olea 2009). The rubbish dump may have benefitted floaters and immature birds more than territorial breeding pairs (Oro *et al.* 2008, Weiser and Powell 2011, Lieury *et al.* 2015), and the closure of the dump may therefore alleviate interference competition by reducing the number of floaters in the study area, which could explain why we did not observe a decrease in breeding success after the closure of the rubbish dump. Alternatively, high plasticity in diet choice, which has been found in Egyptian Vultures (Dobrev *et al.* 2016), may allow adult birds to rapidly shift to other food sources and therefore compensate for the loss of a food source such as the communal rubbish dump.

Our estimates of fledging rate are similar to the range of European populations of Egyptian Vultures (Mateo-Tomás *et al.* 2010), and we also found no effect of the closure of the rubbish dump on fledging rate. The main threat to Egyptian Vultures in Turkey is seen as food shortages arising from declining traditional animal husbandry (Iñigo *et al.* 2008), but in our study area there are still a large number of small agricultural units that can provide food for vultures (Figure 1). These small agricultural units likely provide a temporally heterogeneous supply of potential food sources, and therefore replicate the less predictable opportunistic foraging opportunities to which vultures are adapted (Cortés-Avizanda *et al.* 2010). As a result of these other foraging opportunities, the rubbish dump may not have played a critical role in food acquisition for territorial adult vultures, explaining why we detected no effect of the rubbish dump closure on fledging rate. Maintaining the heterogeneous landscape mosaic of small animal husbandry operations that can provide small amounts of food at unpredictable times may therefore be more beneficial to the Egyptian Vulture population than large and permanent central feeding stations such as rubbish dumps or vulture restaurants due to reduced effects of interference competition (Cortés-Avizanda *et al.* 2012). However, some of the chicken farm disposal sites that are promoted as grassland fertilization appear to be industrial waste disposal sites in the open landscape. If the chickens reared in those industrial chicken farms that dispose waste material on grassland are treated with pharmaceuticals that can be harmful to vultures, then survival probabilities of adult vultures may be negatively affected despite short-term improvements in fledging rate (Green *et al.* 2006, Ogada *et al.* 2012). We recommend that animal products made available for vultures in the open landscape are tested for substances that are known to be harmful to vultures, and that problematic pharmaceuticals are banned for veterinary use in Turkey (Margalida *et al.* 2014a).

In summary, we conclude that the closure of the rubbish dump in Beypazarı has not led to a substantial increase or decrease in reproductive parameters, but we recommend continued monitoring as the effects of a lower floater population may potentially improve breeding success over time. In addition, because feeding stations are known to benefit the survival of immature birds (Oro *et al.* 2008, Margalida *et al.* 2014b, Lieury *et al.* 2015), a better understanding of other demographic parameters such as adult and juvenile survival and recruitment is required to assess whether the population is sustainable with the current levels of productivity.

## Acknowledgements

We acknowledge the support of colleagues helping to monitor territories, especially Adem Akyol, Mustafa Akyol, Fatih Bük, Şenol Uzunoğlu, İzzet Koçak, Osman Türkdöğün, Bahattin Özcan, Emrah Varol, Onur Güngör, and Mehmet Ertuğrul. This work was financially supported by the LIFE+ project “The Return of the Neophron” (LIFE10 NAT/BG/000152) funded by the European Union and co-funded by the AG Leventis Foundation. We appreciate constructive comments by Antoni Margalida, Metodija Veleviski, and an anonymous reviewer on an earlier draft of the paper.



## References

- Angelov, I., Hashim, I. and Opper, S. (2013) Persistent electrocution mortality of Egyptian Vultures *Neophron percnopterus* over 28 years in East Africa. *Bird Conserv. Internatn.* 23: 1–6.
- Bates, D., Maechler, M., Bolker, B. and Walker, S. (2015) Fitting Linear Mixed-Effects Models using lme4. *J. Stat. Softw.* 67: 1–48.
- Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H. and White, J.-S. S. (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends Ecol. Evol.* 24: 127–35.
- Bougain, C. (2016) Identification of important migration concentration areas of Egyptian vultures *Neophron percnopterus* from the Balkan population tracked by satellite telemetry. MSc thesis, Strasbourg, France: University of Strasbourg.
- Cameron, R. A. D., Cornwallis, L., Percival, M. J. L. and Sinclair, A. R. E. (1967) The migration of raptors and storks through the Near East in autumn. *Ibis* 109: 489–501.
- Carrete, M., Donazar, J. A. and Margalida, A. (2006) Density-dependent productivity depression in Pyrenean Bearded Vultures: Implications for conservation. *Ecol. Appl.* 16: 1674–1682.
- Carrete, M., Grande, J. M., Tella, J. L., Sánchez-Zapata, J. A., Donazar, J. A., Díaz-Delgado, R. and Romo, A. (2007) Habitat, human pressure, and social behavior: Partialling out factors affecting large-scale territory extinction in an endangered vulture. *Biol. Conserv.* 136: 143–154.
- Ceballos, O. and Donazar, J. A. (1989) Factors influencing the breeding density and nest-site selection of the Egyptian Vulture (*Neophron percnopterus*). *J. Ornithol.* 130: 353–359.
- Chaudhary, A., Subedi, T. R., Giri, J. B., Baral, H. S., Bidari, B., Subedi, H., Chaudhary, B., Chaudhary, I., Paudel, K. and Cuthbert, R. J. (2012) Population trends of critically endangered *Gyps* vultures in the lowlands of Nepal. *Bird Conserv. Internatn.* 22: 270–278.
- Cortés-Avizanda, A., Carrete, M. and Donazar, J. A. (2010) Managing supplementary feeding for avian scavengers: Guidelines for optimal design using ecological criteria. *Biol. Conserv.* 143: 1707–1715.
- Cortés-Avizanda, A., Jovani, R., Carrete, M. and Donazar, J. A. (2012) Resource unpredictability promotes species diversity and coexistence in an avian scavenger guild: a field experiment. *Ecology* 93: 2570–2579.
- Cuthbert, R., Green, R. E., Ranade, S., Saravanan, S., Pain, D. J., Prakash, V. and Cunningham, A. A. (2006) Rapid population declines of Egyptian vulture (*Neophron percnopterus*) and red-headed vulture (*Sarcogyps calvus*) in India. *Anim. Conserv.* 9: 349–354.
- Demerdzhiev, D., Stoychev, S., Dobrev, D., Spasov, S. and Opper, S. (2015) Studying the demographic drivers of an increasing Imperial Eagle population to inform conservation management. *Biodivers. Conserv.* 24: 627–639.
- Dobrev, V., Boev, Z., Arkumarev, V., Dobrev, D., Kret, E., Saravia, V., Bounas, A., Vavylis, D., Nikolov, S. C. and Opper, S. (2016) Diet is not related to productivity but to territory occupancy in a declining population of Egyptian Vultures *Neophron percnopterus*. *Bird Conserv. Internatn.* 26: 273–285.
- Donald, P. F., Round, P. D., Dai We Aung, T., Grindley, M., Steinmetz, R., Shwe, N. M. and Buchanan, G. M. (2015) Social reform and a growing crisis for southern Myanmar's unique forests. *Conserv. Biol.* 29: 1485–1488.
- Donald, P. F., Sanderson, F. J., Burfield, I. J. and van Bommel, F. P. J. (2006) Further evidence of continent-wide impacts of agricultural intensification on European farmland birds, 1990–2000. *Agric. Ecosyst. Environ.* 116: 189–196.
- Donazar, J. A., Cortés-Avizanda, A. and Carrete, M. (2010) Dietary shifts in two vultures after the demise of supplementary feeding stations: consequences of the EU sanitary legislation. *Eur. J. Wildl. Res.* 56: 613–621.
- Donazar, J. A., Margalida, A., Carrete, M. and Sanchez-Zapata, J. A. (2009) Too sanitary for vultures. *Science* 326: 664.
- Eken, G., Bozdoğan, M., İsfendiyaroğlu, S., Kılıç, D. T. and Lise, Y. (eds) (2006) *Türkiye'nin*

- Önemli Doğa Alanları, Ankara: Doğa Derneği.
- Fielding, A. H. and Bell, J. F. (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environ. Conserv.* 24: 38–49.
- García-Ripollés, C. and López-López, P. (2006) Population size and breeding performance of Egyptian Vultures (*Neophron percnopterus*) in eastern Iberian Peninsula. *J. Raptor Res.* 40: 217–221.
- Gillies, C. S., Hebblewhite, M., Nielsen, S. E., Krawchuk, M. A., Aldridge, C. L., Frair, J. L., Saher, D. J., Stevens, C. E. and Jerde, C. L. (2006) Application of random effects to the study of resource selection by animals. *J. Anim. Ecol.* 75: 887–898.
- González, L. M., Margalida, A., Sánchez, R. and Oria, J. (2006) Supplementary feeding as an effective tool for improving breeding success in the Spanish imperial eagle (*Aquila adalberti*). *Biol. Conserv.* 129: 477–486.
- Green, R. E., Taggart, M. A., Das, D., Pain, D. J., Sashi Kumar, C., Cunningham, A. A. and Cuthbert, R. (2006) Collapse of Asian vulture populations: risk of mortality from residues of the veterinary drug diclofenac in carcasses of treated cattle. *J. Appl. Ecol.* 43: 949–956.
- Grubač, B., Veleviski, M. and Avukatov, V. (2014) Long-term population decrease and recent breeding performance of the Egyptian Vulture *Neophron percnopterus* in Macedonia. *North-West. J. Zool.* 10: 25–32.
- Hazler, K. R. (2004) Mayfield logistic regression: a practical approach for analysis of nest survival. *Auk* 121: 707–716.
- Iñigo, A., Barov, B., Orhun, C. and Gallo-Orsi, U. (2008) *Action plan for the Egyptian Vulture Neophron percnopterus in the European Union*, Brussels: BirdLife International and European Commission.
- Jiménez-Valverde, A. (2012) Insights into the area under the receiver operating characteristic curve (AUC) as a discrimination measure in species distribution modelling. *Glob. Ecol. Biogeogr.* 21: 498–507.
- Kamp, J., Oppel, S., Ananin, A. A., Durnev, Y. A., Gashev, S. N., Hölzel, N., Mishchenko, A. L., Pessa, J., Smirenski, S. M., Strelnikov, E. G., Timonen, S., Wolanska, K. and Chan, S. (2015) Global population collapse in a superabundant migratory bird and illegal trapping in China. *Conserv. Biol.* 29: 1684–1694.
- Kemp, M. U., Emiel van Loon, E., Shamoun-Baranes, J. and Bouten, W. (2012) RNCEP: Global weather and climate data at your fingertips. *Methods Ecol. Evol.* 3: 65–70.
- Kirwan, G. M., Boyla, K. A., Castell, P., Demirci, B., Özen, M., Welch, H. and Marlow, T. (2008) *The birds of Turkey*. London, UK: Christopher Helm.
- Lewis, F., Butler, A. and Gilbert, L. (2011) A unified approach to model selection using the likelihood ratio test. *Methods Ecol. Evol.* 2: 155–162.
- Liberatori, F. and Penteriani, V. (2001) A long-term analysis of the declining population of the Egyptian vulture in the Italian peninsula: Distribution, habitat preference, productivity and conservation implications. *Biol. Conserv.* 101: 381–389.
- Lieury, N., Gallardo, M., Ponchon, C., Besnard, A. and Millon, A. (2015) Relative contribution of local demography and immigration in the recovery of a geographically-isolated population of the endangered Egyptian vulture. *Biol. Conserv.* 191: 349–356.
- López-López, P., García-Ripollés, C. and Urios, V. (2014) Food predictability determines space use of endangered vultures: implications for management of supplementary feeding. *Ecol. Appl.* 24: 938–949.
- Margalida, A. and Colomer, M. A. (2012) Modelling the effects of sanitary policies on European vulture conservation. *Sci. Rep.* 2: 1–7.
- Margalida, A., Benítez, J. R., Sánchez-Zapata, J. A., Ávila, E., Arenas, R. and Donazar, J. A. (2012) Long-term relationship between diet breadth and breeding success in a declining population of Egyptian Vultures *Neophron percnopterus*. *Ibis* 154: 184–188.
- Margalida, A., Bogliani, G., Bowden, C. G. R., Donazar, J. A., Genero, F., Gilbert, M., Karesh, W. B., Kock, R., Lubroth, J., Manteca, X., Naidoo, V., Neimanis, A., Sanchez-Zapata, J. A., Taggart, M. A., Vaarten, J., Yon, L., Kuiken, T. and Green, R. E. (2014a) One health approach to use of veterinary pharmaceuticals. *Science* 346: 1296–1298.

- Margalida, A., Colomer, M. À and Oro, D. (2014b) Man-induced activities modify demographic parameters in a long-lived species: effects of poisoning and health policies. *Ecol. Appl.* 24: 436–444.
- Margalida, A., Donazar, J. A., Carrete, M. and Sánchez-Zapata, J. A. (2010) Sanitary versus environmental policies: fitting together two pieces of the puzzle of European vulture conservation. *J. Appl. Ecol.* 47: 931–935.
- Mateo-Tomás, P. and Olea, P. P. (2009) Combining scales in habitat models to improve conservation planning in an endangered vulture. *Acta Oecologica* 35: 489–498.
- Mateo-Tomás, P. and Olea, P. P. (2010) Diagnosing the causes of territory abandonment by the Endangered Egyptian vulture *Neophron percnopterus*: the importance of traditional pastoralism and regional conservation. *Oryx* 44: 424–433.
- Mateo-Tomás, P., Olea, P. P. and Fombellida, I. (2010) Status of the Endangered Egyptian vulture *Neophron percnopterus* in the Cantabrian Mountains, Spain, and assessment of threats. *Oryx* 44: 434–440.
- Ogada, D. L., Keesing, F. and Virani, M. Z. (2012) Dropping dead: causes and consequences of vulture population declines worldwide. *Ann. N. Y. Acad. Sci.* 1249: 57–71.
- Ogada, D., Shaw, P., Beyers, R. L., Buij, R., Murn, C., Thiollay, J. M., Beale, C. M., Holdo, R. M., Pomeroy, D., Baker, N., Krüger, S. C., Botha, A., Virani, M. Z., Monadjem, A. and Sinclair, A. R. E. (2016) Another continental vulture crisis: Africa's cultures collapsing toward extinction. *Conserv. Lett.* 9: 89–97.
- Oppel, S., Dobrev, V., Arkumarev, V., Saravia, V., Bounas, A., Kret, E., Veleviski, M., Stoychev, S. and Nikolov, S. C. (2015) High juvenile mortality during migration in a declining population of a long-distance migratory raptor. *Ibis* 157: 545–557.
- Oppel, S., Dobrev, V., Arkumarev, V., Saravia, V., Bounas, A., Manolopoulos, A., Kret, E., Veleviski, M., Popgeorgiev, G. S. and Nikolov, S. C. (2017) Landscape factors affecting territory occupancy and breeding success of Egyptian Vultures on the Balkan Peninsula. *J. Ornithol.* 158: 443–457.
- Oppel, S., Iankov, P., Mumun, S., Gerdzhirov, G., Iliev, M., Isfendiyaroglu, S., Yenyurt, C. and Tabur, E. (2014) Identification of the best sites around the gulf of Iskenderun, Turkey, for monitoring the autumn migration of Egyptian Vultures and other diurnal raptors. *Sandgrouse* 36: 240–249.
- Oro, D., Margalida, A., Carrete, M., Heredia, R. and Donazar, J. A. (2008) Testing the goodness of supplementary feeding to enhance population viability in an endangered vulture. *PLoS One* 3: e4084.
- Porter, R. and Willis, I. (1968) The autumn migration of soaring birds at the Bosphorus. *Ibis* 110: 520–536.
- R Core Team (2016) *R: A language and environment for statistical computing*, Vienna, Austria: R Foundation for Statistical Computing.
- Sanz-Aguilar, A., Sánchez-Zapata, J. A., Carrete, M., Benítez, J. R., Ávila, E., Arenas, R. and Donazar, J. A. (2015) Action on multiple fronts, illegal poisoning and wind farm planning, is required to reverse the decline of the Egyptian vulture in southern Spain. *Biol. Conserv.* 187: 10–18.
- Sarà, M. and Vittorio, M. (2003) Factors influencing the distribution, abundance and nest-site selection of an endangered Egyptian vulture (*Neophron percnopterus*) population in Sicily. *Anim. Conserv.* 6: 317–328.
- Şekercioğlu, Ç. H., Anderson, S., Akçay, E., Bilgin, R., Can, Ö. E., Semiz, G., Tavşanoğlu, Ç., Yokeş, M. B., Soyumert, A., İpekdal, K., Sağlam, İ. K., Yücel, M. and Nüzhet Dalfes, H. (2011) Turkey's globally important biodiversity in crisis. *Biol. Conserv.* 144: 2752–2769.
- Şen, B., Tavares, J. P. and Bilgin, C. C. (2017) Nest site selection patterns of a local Egyptian Vulture *Neophron percnopterus* population in Turkey. *Bird Conserv. Internatn.* in press.
- Sutherland, W. J. and Brooks, D. J. (1981) The autumn migration of raptors, storks, pelicans and spoonbills at the Belen Pass, southern Turkey. *Sandgrouse* 2: 1–21.
- Tauler-Ametller, H., Hernández-Matías, A., Pretus, J. L. L. and Real, J. (2017) Landfills determine the distribution of an expanding breeding population of the endangered Egyptian Vulture *Neophron percnopterus*. *Ibis*. in press.
- Tauler, H., Real, J., Hernández-Matías, A., Aymerich, P., Baucells, J., Martorell, C.

- and Santandreu, J. (2015) Identifying key demographic parameters for the viability of a growing population of the endangered Egyptian Vulture *Neophron percnopterus*. *Bird Conserv. Internatn.* 25: 1–14.
- Thiollay, J. M. (2006) The decline of raptors in West Africa: long-term assessment and the role of protected areas. *Ibis* 148: 240–254.
- Velevski, M., Nikolov, S. C., Hallmann, B., Dobrev, V., Sidiropoulos, L., Saravia, V., Tsiakiris, R., Arkumarev, V., Galanaki, A., Kominos, T., Stara, K., Kret, E., Grubač, B., Lisičanec, E., Kastritis, T., Vavylis, D., Topi, M., Hoxha, B. and Opper, S. (2015) Population decline and range contraction of the Egyptian Vulture *Neophron percnopterus* in the Balkan Peninsula. *Bird Conserv. Internatn.* 25: 440–450.
- Virani, M. Z., Kendall, C., Njoroge, P. and Thomsett, S. (2011) Major declines in the abundance of vultures and other scavenging raptors in and around the Masai Mara ecosystem, Kenya. *Biol. Conserv.* 144: 746–752.
- Vlachos, C. G., Papageorgiou, N. K. and Bakalaoudis, D. E. (1998) Effects of the feeding station establishment on the Egyptian Vulture (*Neophron percnopterus*) in Dadia Forest, North Eastern Greece. Pp. 197–207 in R. D. Chancellor B.-U. Meyburg and J. J. Ferrero, eds. *Holarctic birds of prey*. Adenex – World Working Group on Birds of Prey and Owls.
- Weiser, E. L. and Powell, A. N. (2011) Reduction of garbage in the diet of nonbreeding glaucous gulls corresponding to a change in waste management. *Arctic* 64: 220–226.

JAKOB KATZENBERGER

Workgroup on Endangered Species, J.F. Blumenbach Institute of Zoology and Anthropology,  
Georg-August University of Göttingen, Bürgerstrasse 50, 37073 Göttingen, Germany.

EVİRİM TABUR, BİLGEÇAN ŞEN, SÜREYYA İŞFENDİYAROĞLU, İTRİ LEVENT ERKOL  
Doğa Derneği, Kızılay Mah. Menekşe 2 Sok. 33/5 Çankaya/Ankara, Turkey.

STEFFEN OPPEL\*

RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, The David  
Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, United Kingdom.

\*Author for correspondence; e-mail: steffen.oppel@rspb.org.uk

Received 23 April 2017; revision accepted 21 July 2017